

Comparison of Treatment Outcomes in Childhood Pneumonia with or without Rickets

Noor ul Huda¹, Masood Mazhar Sahu², Muhammad Hanif Memon³, Tasneem Kousar⁴

¹Postgraduate resident, Pediatric Medicine department, Reccep Tayyap Erdogan Hospital Muzaffargarh

²Senior Consultant, Pediatric Medicine department, Reccep Tayyap Erdogan Hospital Muzaffargarh

³Assistant Professor of Pediatric department, Suleman Roshan Medical College, Tando Adam

⁴Associate Professor of Pediatric department Suleman Roshan Medical College, Tando Adam

Author's Contribution

¹Substantial contributions to the conception or design of the work; or the acquisition, ²Final Approval of the study to be published ^{3,4}Drafting the work or revising it critically for important intellectual content,

Funding Source: None

Conflict of Interest: None

Received: June 10, 2025

Revised: Dec 02, 2025

Accepted: Jan 09, 2026

Address of Correspondent

Dr. Noor ul Huda

Post graduate resident, Pediatric

Medicine department, Reccep

Tayyap Erdogan Hospital

Muzaffargarh

noorulhuda19111997@gmail.com

ABSTRACT

Objective: To compare the treatment outcomes among children with pneumonia with or without rickets presentation.

Methodology: This comparative study was conducted on 190 children diagnosed with pneumonia, who were divided into two equal groups: 95 children with rickets and 95 without rickets. Study was done at Department of Pediatric Medicine, Recep Tayyip Erdogan Hospital Muzaffargarh. The rickets was diagnosed based on clinical features and radiological and biochemical findings. Patients received standard pneumonia management according to institutional guidelines. Treatment outcomes were assessed in terms of duration of fever, need for oxygen therapy, length of hospital stay, and treatment failure, across the groups. Data were analyzed using appropriate statistical tests, and a *p*-value of < 0.05 was considered statistically significant.

Results: Paediatric patients with rickets showed a significantly longer duration of fever (*p*=0.001) and prolonged stay in Hospital stay *p*=0.001 compared to non-rickets patients. Failure of treatment was more frequent in the rickets group (22.1 % vs 9.5 %) *p* = 0.018. The severe pneumonia and oxygen requirement were significantly higher in children with rickets in contrast to non-rickets patients (31.6 % vs 15.8 %) and (48.4 % vs 27.4 %), *p*= 0.004 respectively. Average clinical recovery time was also significantly delayed in children with rickets. Additionally, the rate of complications also being significantly higher among children with rickets, while the mortality rate was low and almost similar across the *p*=0.05). **Conclusion:** The childhood pneumonia with rickets was observed with significantly poor treatment outcomes, and these patients required more intensive supportive care and showed delayed recovery compared to those without rickets.

Key words: Pneumonia, Rickets, Children, Treatment failure, Mortality.

Cite this article as: Huda NU, Sahy MM, Memon MH, Kousar T. Comparison of Treatment Outcomes in Childhood Pneumonia with or without Rickets. *Ann Pak Inst Med Sci.* 2026; 22(1):14-19. doi. 10.48036/apims.v22i1.1545

Introduction

Pneumonia is globally recognized as one of the leading causes of childhood mortality, especially among children between 6 to 60 months of age worldwide.¹ It ranks as the 2nd most common cause of mortality in this age group, responsible for more than 30,000 deaths per year.² Regardless advances in antibiotic therapy and vaccination coverage, failure of treatment and prolonged hospital stays continue to pose a major challenges clinically, specifically in resource-limited settings where underlying nutritional deficiencies are predominant.

The Rickets, is a metabolic bone disease produced mainly by deficiency of vitamin D, decreased level of calcium, or phosphate, remains a considerable public health concern in developing populations, mostly characterized by widening and flaring of the metaphysis of bones, subsequent in enlargement of the wrist joints, prominence of the costochondral junctions of the ribs, knock-knees, and bowed legs.³ Middle Eastern, African and Asian countries report an extensive prevalence of rickets among child population with prevalence of it 15-18% in the South Asia and 10% in the Africa.³ The risk factors for developing vitamin D deficiency rickets incorporate

inadequate exposure to sunlight, exclusive breastfeeding without supplementation of vitamin D and insufficient dietary intake of vitamin D, phosphorus or the calcium. In Pakistan and similar low-income settings, these risk factors are widely prevalent due to poverty, malnutrition, and limited access to preventive healthcare facilities.⁴

The diagnosis of nutritional rickets is primarily made through clinical assessment, including the medical history and physical examination of the patients, further supported by biochemical investigations and ultimately verified through radiographic parameters.⁵ However, the nutritional rickets can lead to several health complications, like decreased linear growth, chest wall deformities that raise the susceptibility to pneumonia, fractures, pain of the bones, and lower limbs deformities. Moreover, affected children may happening delays in gross motor development, and in cases with severe condition, life-threatening complications may occur, including hypocalcemia related seizures, and cardiovascular events.⁵⁻⁷

The relationship between rickets and pneumonia is of particular clinical importance and the treatment outcomes has been explored in a limited number of studies. The epidemiological studies have decisively established the link between nutritional rickets and the risk of pneumonia development, as according to a study conducted in Ethiopia reported that, there was a 13-fold greater incidence of rickets among children with pneumonia than among controls, indicating that the deficiency of vitamin D or calcium may be important predisposing factors for pneumonia in children under 5 years in developing nations.⁸

In the same way, the Nepalian study measured plasma 25(OH)D concentrations among 568 children hospitalized with WHO-defined severe pneumonia and found that plasma 25(OH)D levels below 50 nmol/L were corelated with the raised risk of treatment failure and prolonged duration for recovery.⁹ Such previous findings underscore the potential of rickets as an important modifiable risk factor influencing clinical outcomes in pneumonia among children. In spite of this growing evidence base, comparative data on treatment outcomes specifically between rachitic and non-rachitic children with pneumonia remain very limited, specifically from South Asian settings including Pakistan where both conditions are highly frequent. However present study was therefore planned to compare treatment outcomes, including treatment failure, time to fever resolution, normalization of respiratory rate, and duration of hospital stay, among

children with pneumonia with and without co-existing rickets, in order to produce locally relevant evidence that may inform clinical management and highlight the nutritional screening importance among children hospitalized with pneumonia.

Methodology

A prospective comparative cohort study was conducted in the Department of Pediatric Medicine at Recep Tayyip Erdogan Hospital from January 2025 to June 2025. Ethical approval was obtained from the hospital's ethics committee (Ref. No. IHHN_IRB_2024_12_003).

All children aged 2 to 12 years, of either gender, presenting with pneumonia as per the operational definition were included. Pneumonia was defined as the presence of cough, fever (axillary temperature $>100^{\circ}\text{F}$), and increased respiratory rate (>50 breaths/min in children aged 2–11 months, >40 in 2–5 years, and >30 in children older than 5 years), with a duration of ≤ 72 hours. Children with and without rickets were included.

Exclusion criteria included children with chronic lung diseases (such as asthma, cystic fibrosis, or post-tuberculosis bronchiectasis based on history and medical record review), immunosuppressed children, and those whose parents did not consent to participate.

Children were enrolled after obtaining informed consent from their parents. Baseline data—including age, gender, history of exclusive breastfeeding, nutritional status, and vaccination status—were recorded. A detailed clinical examination was performed in all participants to identify bony manifestations of rickets. Additionally, 3 mL of venous blood was collected from each child for biochemical analysis.

Rickets was diagnosed based on the operational definition: the presence of one or more bony abnormalities (such as bowing of the legs, knock knees, rachitic rosary, swelling of the ends of long bones, or frontal bossing) along with biochemical abnormalities, including serum 25-hydroxyvitamin D levels <20 ng/mL, serum calcium <8.5 mg/dL, low phosphorus levels, elevated alkaline phosphatase (>350 IU/L), and serum phosphate <4.0 mg/dL.

Participants were categorized into two groups: the exposed group (rickets-positive) and the unexposed group (without rickets). All children received standard management, including oxygen therapy, intravenous antibiotics, and treatment for rickets (where applicable), according to institutional protocols.

Patients were followed until their final outcomes. Treatment failure was defined as the absence of clinical improvement within 48–72 hours of initiating standard antibiotic therapy, indicated by persistent fever (>38.5°C), ongoing respiratory distress, or oxygen saturation below 95%, necessitating a change in antibiotics or admission to the intensive care unit.

Additionally, time to fever resolution, normalization of respiratory rate, and duration of hospital stay were recorded. All data were entered into a structured proforma and analyzed using SPSS version 23.

Results

This study enrolled 100 patients, which were further equally divided into two groups; 50 in each group. Overall majority of boys (60%), and most children were not exclusively breastfed (67%). The malnutrition was uncommon overall, affecting only 9% of the total sample, while it was slightly more prevalent in the rickets group (12%) compared to the non-rickets group (6%). According to the vaccination status, around half of all children were partially vaccinated (48%) and the other half fully vaccinated (48%), with only 4 unvaccinated children, all from the unexposed group. Overall baseline characteristics, were statistically insignificant across the groups (p >0.05). Table I

In this study the treatment failure was noted higher in cases of rickets group 8% compared to the non-rickets group (2%). Moreover, exposed group approximately 4 times more likely to have treatment failure than those without rickets (OR 4.261, 95% CI: 0.459–39.544), indicating the

presence of rickets is associated with higher risk of poor treatment response in childhood pneumonia, while this association did not reach statistical significance given the wide CI, possibly due to the very limited number of failure events. Figure 1.




Figure 1. Frequency of treatment failure in both groups. (n=100)

Overall the children presented with rickets had significantly worse recovery outcomes compared to those without rickets, particularly as the fever took more than twice as long to resolve in the rickets group (6.32 days) compared to the non-rickets group (3.12 days) (p=0.001). Likewise, normalization of respiratory rate was also significantly delayed in rickets group (8.48 days) in contrast to children without rickets (3.20 days) (p=0.001). Additionally, the mean Hospital stay was also noted longer in the rickets group (12.24 days) compared to the non-rickets group (5.88 days) (p=0.001), as presented in table: II.

Table I: Baseline and clinical characteristics of patients in both groups. (n=100)

VARIABLES	STUDY GROUPS		Total	p-value
	Exposed (Rickets positive)	Unexposed (Without rickets)		
Gender	Boys	32 64.0%	28 56.0%	0.309
	Girls	18 36.0%	22 44.0%	
Exclusive breastfeeding	Yes	16 32.0%	17 34.0%	0.529
	No	34 68.0%	33 66.0%	
Malnutrition	Yes	6 12.0%	3 6.0%	0.505
	No	44 88.0%	47 94.0%	
Vaccinated status	Unvaccinated	0 0.0%	4 8.0%	0.063
	Partially vaccinated	26 52.0%	22 44.0%	
	Fully vaccinated	24 48.0%	24 48.0%	

Table II: Comparison of recovery outcomes in both groups. (n=100)

Variables	Study groups	N	Mean	SD	p-value
Time to fever resolution (days)	Exposed (Rickets positive)	50	6.32	1.86	0.001
	Unexposed (Without rickets)	50	3.12	1.25	
Time to normalization of respiratory rate (days)	Exposed (Rickets positive)	50	8.48	3.58	0.001
	Unexposed (Without rickets)	50	3.20	1.61	
Duration of Hospital stay (days)	Exposed (Rickets positive)	50	12.24	4.35	0.001
	Unexposed (Without rickets)	50	5.88	1.68	

Discussion

The nutritional rickets is associated to poorer clinical outcomes and a prolonged disease course in children with pneumonia, often demanding repeated changes in antibiotic therapy. Both conditions are highly frequent in our population and are main contributors to childhood morbidity; though, there is very limited data evaluating how rickets influences the clinical progression and recovery of pneumonia among children.¹⁰ This study was comparatively conducted to evaluate the impact of rickets on the course of pneumonia by comparing important clinical features and symptom resolution between pediatric patients with and without rickets, and there were boys in majority 60% compared to girls 41% and this gender analysis was almost similar to the study by Jamal A et al¹⁰ where male patients were 68 (56.1%) and female patients were 53 (43.8%).

In this study the treatment failure rate was significantly higher 8% in the rickets group compared to 2% only in non-rickets group, representing a fourfold difference. These findings were aligned with few other investigations, like Banajeh et al¹¹ conducted a prospective cohort study on 152 children with WHO-defined very severe pneumonia, stated that treatment failure occurred in 24 children overall, with 21 of those (20.6%) being rachitic, which was significantly higher among in rachitic children compared to the non-rachitic children (6%) (OR 1.38, $P=0.031$). In the study by Jakhar SK et al¹² also reported that the children with sign of rickets had significantly higher rate of treatment failure 7 (47%) ($p=0.054$), prolonged Hospital stay 18 (36%) ($p=0.019$) and need for change in antibiotics 15 (44%) ($p=0.002$). Additionally, a Nepalese study measuring plasma 25(OH)D in 568 children hospitalized with severe pneumonia found that vitamin D levels below 50 nmol/L were independently correlated with raised risk for treatment failure and prolonged recovery duration, significantly associating subclinical vitamin D insufficiency often the precursor to clinical rickets with the adverse outcomes.¹³ Furthermore in the study by Jamal A et al¹⁰ observed that the Hospitalized children with pneumonia and rickets

identified with prolonged and significantly delayed clinical recovery, specifically in resolution of fever, tachypnea, cough, and return to normal feeding in contrast to those without rickets.¹⁰ In aligns to our findings Lakshmi M et al¹⁴ demonstrated that the according to their multivariate regression analysis the infancy, incomplete immunization, malnutrition, severity of anemia, clinical signs of rickets, temperature, and baseline hypoxia were the significant predictors of treatment failure, specifically for signs of rickets failure rate was 22 (51.16%) compared to 41 (21.40%) among those without signs of rickets ($p=0.015$). The delayed recovery from pneumonia and higher treatment failure rate among children with rickets may because the rickets and its underlying vitamin-D deficiency impair immune function, oxygenation, and overall flexibility in nutritional status, leading to sluggish clearance of infection and more complications.

Furthermore, in this study the rickets group showed significantly longer duration to achieve fever resolution (mean 6.32 ± 1.86 days vs. 3.12 ± 1.25 days), normalization of respiratory rate (mean 8.48 ± 3.58 days vs. 3.20 ± 1.61 days) and hospitalization (mean 12.24 ± 4.35 days vs. 5.88 ± 1.68 days), in rickets group versus non-rickets group ($p=0.001$). In aligns to this series, the Jamal A et al¹⁰ in their cohort of 121 hospitalized children aged 1–5 years with pneumonia, reported that, those with rickets experienced a significantly prolonged time to clinical stabilization compared to children without rickets.

Jamal A et al.¹⁵ reported that children with pneumonia and coexisting rickets require a significantly longer time to achieve normalization of respiratory rate compared to those with pneumonia alone. Similarly, Haugen J et al.¹³ observed that children with lower vitamin D levels (25(OH)D <50 nmol/L) had higher rates of treatment failure and experienced delayed recovery in cases of severe lower respiratory tract infections.

In addition, Piloya et al.¹⁶, in their cross-sectional study of 221 children aged 2–59 months with severe pneumonia, reported a 9.5% prevalence of nutritional rickets. They highlighted that rickets is a common comorbidity in children admitted with pneumonia, even in tropical regions with abundant sunlight, and recommended routine

screening for rickets in all children presenting with severe pneumonia.

In this study the exclusive breastfeeding was found in 32% of the group and 34% of the non-rickets group, and malnutrition was identified among 12% of children with rickets contrasted with 6% of children without rickets. Such patterns were closely parallel with other studies like Altaib RM et al¹⁷ reported that the 62% of the participating children exhibited signs of rickets along with low level of vitamin D. Particularly, exclusive breastfeeding during the early months of life was reported in 74% of the children. Comparable trends have been studied in other studies; where reported that the most of the of children who developed rickets were exclusively breastfed.¹⁸⁻¹⁹ Overall in the several low-resource countries like ours, the exclusive breastfeeding is very common and often prolonged without vitamin-D supplementation, which significantly raise the risk of nutritional rickets among infants and young children. However, the condition, in turn, is significantly linked to delayed recovery and higher treatment failure among children with pneumonia, may because the vitamin-D deficiency impairs immunity, worsens hypoxemia, and decreases overall nutritional replacement. Overall the substantial proportion of rachitic children in recent studies are exclusively breastfed,^{16,18,20,21} ensuring routine vitamin-D supplementation from early infancy is very important and cost-effective measure to prevent rickets and thereby improve pneumonia outcomes in the children. This study has several limitations, including a relatively limited sample size and a single-center design, which may affect generalizability and increases the risk of selection bias. Consequently, it is strongly recommended that large-scale, multicenter studies be conducted to confirm the findings and more precisely quantify the impact of rickets on pneumonia outcomes among children.

Conclusion

The children with rickets and parallel pneumonia observed with significantly higher treatment failure rate and longer recovery, in terms of prolonged fever, delayed respiratory recovery, and extended hospitalization in contrast to children without rickets underscoring the adverse impact of rickets on pneumonia outcomes. Generally, findings highlighted the need for early screening and correction of vitamin-D deficiency and rickets, along with routine vitamin-D supplementation, to decrease the failure rate of treatment and improve recovery among children with pneumonia.

References

1. Bantie GM, Meseret Z, Bedimo M, Bitew A. The prevalence and root causes of delay in seeking healthcare among mothers of under five children with pneumonia in hospitals of Bahir Dar city, North West Ethiopia. *BMC Pediatr*. 2019;19(1):482. <https://doi.org/10.1186/s12887-019-1869-9>
2. Kumar A, Chandio K, Mirjat S, Malhi MM, Sathio S, Rahimoon RU, Hafeez M. Exploring the risk factors regarding pneumonia among children under the age of five years. *Front Med Health Res*. 2025;3(4):147-53.
3. Mumtaz A, Akram J, Nazir N, Hasan AH, Ali R, Basharat A, Tauqeer A, Warraich JI, Warraich BI, Khan TM. Risk factors of nutritional rickets among children under-five years of age. *Common Health Cond Dis Their Effects Hum Beings*. 2022;7(10):14-19. <https://doi.org/10.47672/ajhmn.1157>
4. Riaz H, Finlayson AE, Bashir S, Hussain S, Mahmood S, Malik F, Godman B. Prevalence of vitamin D deficiency in Pakistan and implications for the future. *Expert Rev Clin Pharmacol*. 2016;9(2):329-38. <https://doi.org/10.1586/17512433.2016.1122519>
5. Roth DE, Abrams SA, Aloia J, Bergeron G, Bourassa MW, Brown KH, Calvo MS, Cashman KD, Combs G, De-Regil LM, Jefferds ME. Global prevalence and disease burden of vitamin D deficiency: a roadmap for action in low- and middle-income countries. *Ann N Y Acad Sci*. 2018;1430(1):44-79. <https://doi.org/10.1111/nyas.13968>
6. Basatemur E, Sutcliffe A. Incidence of hypocalcemic seizures due to vitamin D deficiency in children in the United Kingdom and Ireland. *J Clin Endocrinol Metab*. 2015;100(1):E91-5. <https://doi.org/10.1210/jc.2014-2773>
7. Chehade H, Girardin E, Rosato L, Cachat F, Cotting J, Perez MH. Acute life-threatening presentation of vitamin D deficiency rickets. *J Clin Endocrinol Metab*. 2011;96(9):2681-3. <https://doi.org/10.1210/jc.2011-1112>
8. Muhe L, Lulseged S, Mason KE, Simoes EA. Case-control study of the role of nutritional rickets in the risk of developing pneumonia in Ethiopian children. *Lancet*. 1997;349(9068):1801-4. [https://doi.org/10.1016/S0140-6736\(96\)12098-5](https://doi.org/10.1016/S0140-6736(96)12098-5)
9. Haugen J, Basnet S, Hardang IM, Sharma A, Mathisen M, Shrestha P, Valentiner-Branth P, Strand TA. Vitamin D status is associated with treatment failure and duration of illness in Nepalese children with severe pneumonia. *Pediatr Res*. 2017;82(6):986-93. <https://doi.org/10.1038/pr.2017.71>
10. Jamal A, Tanzil S, Yousuf Yahya SN. Role of rickets in recovery from pneumonia in pediatric inpatient population at a tertiary care hospital in Karachi. *Pak Pediatr J*. 2023;47(1):22-7.
11. Banajeh SM. Nutritional rickets and vitamin D deficiency-association with the outcomes of childhood very severe pneumonia: a prospective cohort study. *Pediatr Pulmonol*. 2009;44(12):1207-15. <https://doi.org/10.1002/ppul.21121>

12. Jakhar SK, Pandey M, Shah D, Ramachandran VG, Saha R, Gupta N, Gupta P. Etiology and risk factors determining poor outcome of severe pneumonia in under-five children. *Indian J Pediatr.* 2018;85(1):20-4. <https://doi.org/10.1007/s12098-017-2514-y>
13. Haugen J, Basnet S, Hardang IM, Sharma A, Mathisen M, Shrestha P, Valentiner-Branth P, Strand TA. Vitamin D status is associated with treatment failure and duration of illness in Nepalese children with severe pneumonia. *Pediatr Res.* 2017;82(6):986-93. <https://doi.org/10.1038/pr.2017.71>
14. Lakshmi M, Manu MA, Kalpan RY, Sabapath S, Rekha TD. Predictors of treatment failure in severe pneumonia treated with ampicillin in tertiary care center. *Int J Contemp Pediatr.* 2019;6:1085-9. <https://doi.org/10.18203/2349-3291.ijcp20191479>
15. Jamal A, Latif I, Aleem SA. Comparison of respiratory rate in children having pneumonia with and without rickets. *Ann Abbasi Shaheed Hosp Kar Med Dent Coll.* 2016;21(4).
16. Piloya T, Odongkara B, Were EM, Ameda F, Mworzi E, Laigong P. Nutritional rickets among children admitted with severe pneumonia at Mulago hospital, Uganda: a cross-sectional study. *BMC Pediatr.* 2018;18(1):336. <https://doi.org/10.1186/s12887-018-1310-9>
17. Altaib RM, Yonis KO, Akrim FA, Qowaider SR, Muhammed AA, Bofarraj M, Ali EM. Nutritional rickets among children admitted with wheezy chest at Al-Bieda Medical Center-Libya. 2021;35(15):70.
18. Khan I, Marwat ZI. Social dilemma and vitamin D deficiency: a cross-sectional study of nutritional rickets in children ages 6 months to 24 months. *Int J Contemp Pediatr.* 2019;6(3):1367-72. <https://doi.org/10.18203/2349-3291.ijcp20192045>
19. Suksantilerd S, Thawatchai R, Rungrojjananon N. Prevalence of vitamin D deficiency in exclusively breastfed infants at Charoenkrung Pracharak Hospital. *World J Clin Pediatr.* 2024;13(1):86693. <https://doi.org/10.5409/wjcp.v13.i1.86693>
20. Santosh S, Arif F, Raess H. Risk factors for nutritional rickets in children under 36 months: a civil hospital case study. *Pak J Med Dent.* 2013;2(3):13-16.
21. Kumaran CRV, Sivakumar KV, Kumar RS. Vitamin D status in exclusively breastfed infants and their mothers. *Int J Contemp Pediatr.* 2021;8:1675-9. <https://doi.org/10.18203/2349-3291.ijcp20213728>